

Rene Birkner

"Incoherent components of the Toric Hilbert scheme"

ZEIT:

19.10.2009, 16:15 Uhr

ORT:

Freie Universitaet Berlin
Institut fuer Mathematik
Arnimallee 3, Rm. 119

The classical Hilbert scheme is the scheme whose closed points are all subschemes of $\mathbb{P}^n$ with the same Hilbert function. That is, for $S = k[x_0, \dots, x_n]$ and an ideal $I \subset S$ the function $H(t) = \dim_k (S/I)_t$ whose value at d is the dimension over k of the degree t part of S/I . Endowing the ring S with a multigrading, i.e. the degree of x_i is $a_i \in \mathbb{N}$, we construct the multigraded Hilbert function. This is the analogon to the classical Hilbert function with degrees in $\mathbb{N}$ for some $d > 0$. Then one can consider all ideals $I \subset S$ with the same multigraded Hilbert function, these are the closed points of the multigraded Hilbert scheme. We consider the simplest case, taking the semigroup $\mathbb{N}^{n+1}$ and the multigraded Hilbert function

$1 \# 1$

Kontakt:

Humboldt-Universität zu Berlin . Institut für Mathematik
SFB 647 . Unter den Linden 6 . 10099 Berlin
Tel. +49 30 2093 1804 . Fax. +49 30 2093 2727
sfb647@math.hu-berlin.de

www.raumzeitmaterie.de