



SFB-Seminar Representation Stability (Teilprojekte C1 u. C3)

ZEIT:

7.1.2014, 16:00 Uhr - 18:00 Uhr

ORT:

Freie Universität Berlin
Zuse-Institut, Hörsaal 2005 EG
Takustraße 7
14195 Berlin-Dahlem

PROGRAMM:

16:00 - 16:45 **Prof. Benson Farb (University of Chicago)**

Representation stability: a user's guide

"Representation stability" refers to a phenomenon discovered a few years ago by Church-Farb that seems to occur all over mathematics; it was developed into a powerful theory with Ellenberg. One simple application gives results such as: the sequence of vector spaces V_n has dimension equal to a polynomial $P(n)$ for n large enough. A common application is to the fixed degree (co)homology of a sequence of spaces X_n .

This has been applied to examples in algebraic topology (configuration spaces), algebraic geometry (moduli spaces of surfaces with n marked points, spaces of polynomials on rank varieties), number theory (cohomology of congruence subgroups), algebraic combinatorics (co-invariant algebras), and several other areas. In most cases nothing is known about the actual dimension of V_n , but this is now reduced in principle to a finite problem. The purpose of this talk will be explain to workers in different areas what this theory can do for them, and how they can apply it.

16:45 - 17:15 Kaffee-Pause

17:15 - 18:00 **Prof. Benson Farb (University of Chicago)**

Kontakt:

Humboldt-Universität zu Berlin . Institut für Mathematik
SFB 647 . Unter den Linden 6 . 10099 Berlin
Tel. +49 30 2093 1804 . Fax. +49 30 2093 2727
sfb647@math.hu-berlin.de

www.raumzeitmaterie.de

Representation stability in cohomology and asymptotics for families of varieties over finite fields

In this talk Prof. Benson Farb will consider two families X_n of varieties on which the symmetric group S_n acts: the configuration space of n points in $\mathbb{C}$ and the space of n linearly independent lines in $\mathbb{C}^n$. He will explain via these two beautiful examples how non-experts can use the (twisted) Grothendieck-Lefschetz Fixed-Point Theorem in ℓ -adic cohomology as a machine to convert information, as follows:

Input: How the multiplicity of a given irreducible representation V of S_n in $H^*(X_n; \mathbb{Q})$ varies with n

Output: Formulas for the number of polynomials over $\mathbb{F}_q$ (resp. maximal tori in $\mathrm{GL}_n(\mathbb{F}_q)$) with specified properties related to V .

In particular we explain how representation stability of $H^*(X_n; \mathbb{Q})$ corresponds to asymptotic stability of various point counts as $n \rightarrow \infty$.

Kontakt:

Humboldt-Universität zu Berlin . Institut für Mathematik
SFB 647 . Unter den Linden 6 . 10099 Berlin
Tel. +49 30 2093 1804 . Fax. +49 30 2093 2727
sfb647@math.hu-berlin.de

www.raumzeitmaterie.de